



Specialists in KEYHOLE Surgery

Incorporating the Australian Centre for Hyperhidrosis & Facial Blushing

Knox Private Hospital, 262 Mountain Highway WANTIRNA VIC 3152
The Valley Private Hospital, Cnr Police & Gladstone Rds, MULGRAVE VIC 3170
Tel: +61 3 9210 7277 Fax: +61 3 9887 5634
Web: www.lapsurgeryaustralia.com.au

Surgeons
Mr Chris Hensman
Mr Ray McHenry
Mr Richard Gilhome

ETS Surgery for Severe Facial Blushing Detailed Information for Prospective Patients.

The following is a detailed account of the effects, complications and side-effects of ETS surgery for severe facial blushing (Idiopathic Craniofacial Erythema). It is based on both our own original research and experience together with scientific information gained at the meeting of the International Society for Sympathetic Surgery held at Erlangen, Germany.

If you are seriously contemplating ETS surgery for facial blushing you must read and familiarise yourself with the following information. This document should be read in conjunction with the more general information provided on our web site at www.lapsurgeryaustralia.com.au. Before proceeding with the operation you must make sure that you understand the entire document and if not that you clarify any issues with your surgeon BEFORE the operation. You will have the opportunity to ask questions of your surgeon and you are encouraged to research both your blushing condition and the treatment options prior to your first consultation. You will be asked to sign a consent form acknowledging that you have read and understood this document and to initial each page of this document to confirm that you have read it.

The Cause of Idiopathic Craniofacial Erythema.

The exact cause is unknown, but it is now classified as a disease of the autonomic (involuntary) nervous system. It is linked to the syndromes of craniofacial, palmar and axillary hyperhidrosis (face, hand and armpit sweating). There is now evidence that it is a genetic condition (autosomal dominant with variable penetration) with up to 50% of people affected having a family history of severe blushing or one of the sweating syndromes.

There is no evidence that it is a primary psychological condition although there can be severe psychological *consequences* from the condition.

Who Should Have the ETS Operation for Severe Facial Blushing?

Surgery for any condition should always be the last resort. Blocking the sympathetic nerves will, for the majority of people with severe facial blushing, provide a significant improvement in quality of life. This assumes that your facial blushing condition is a major impediment to your lifestyle. With any surgery, complications can occur even though they may be unusual.

Side-effects with ETS surgery can occur and a small minority of people, less than 5% in most published studies, will regret having the operation. For around 85 - 90% of people who have the surgery for the right reason, the result will be a significant improvement in quality of life.

Before considering ETS surgery we recommend, if you have not already done so, that you attend your family doctor for a thorough medical examination looking for any other cause of the abnormal blushing. Other possible causes include an overactive thyroid gland, certain other diseases of the endocrine system, some dermatological conditions and certain medications.

Are There Any Alternatives to Surgery?

A variety of non-surgical treatments have been suggested and tried for severe facial blushing. These include medications such as beta blockers, clonidine, antidepressants and a variety of psychological and alternative treatments. Although some individuals may be helped, the majority will gain no significant benefit from these non-surgical treatments. The only treatment which is known scientifically to work for severe facial blushing is ETS surgery.

Nonetheless we recommend that you try some or all of these non-surgical treatments before going ahead with ETS surgery.

The ETS Operation for Severe Facial Blushing.

There is consensus, as confirmed at the fifth meeting of the International Society for Sympathetic Surgery in May 2003, that the correct operation for severe facial blushing is to clip the sympathetic chain at the level of the second rib just below the stellate ganglion. The effects, complications and side-effects of this procedure are discussed later in this document. We now EXCLUSIVELY perform clipping of the nerve rather than cutting the nerve because of the small risk of severe compensatory sweating which is further discussed below under the heading **Side-Effects**. Clipping the nerve is just as effective as cutting it in terms of achieving the desired effect but clipping offers the THEORETICAL chance of reversal in the event of severe compensatory sweating. It must be emphasised that although there are individual reports of successful reversal of the operation by removing the clips, there is as yet no scientific evidence that reversal is possible by removing the clips.

We cannot guarantee reversal of the operation in the event of severe compensatory sweating or other undesired outcome.

The Kuntz Nerve.

If you research the topic of ETS, you will come across various claims and counter-claims about the importance or otherwise of the Kuntz nerve. The Kuntz nerve is a small nerve fibre sometimes seen on the second rib not far from the main sympathetic chain. Its function is not known in humans. Some web-sites on ETS claim success rates of up to 100% for facial blushing because they search for and destroy the Kuntz nerve(s). These same people also claim to be able to correct failed ETS operations by reoperating and destroying the Kuntz nerve.

At the meeting of the International Society for Sympathetic Surgery in Germany, May 2003, attended by a majority of the world's experts in ETS surgery (including us), *all but one of the surgeons present were of the opinion that the Kuntz nerve played no part in the success or failure of ETS surgery for facial blushing. We share this majority opinion.*

Effects.

Effects are the intended outcomes of the operation. For facial blushing, 85-90% of patients will experience either total cessation of blushing or a marked decrease in the severity and frequency of the blushing so as to produce a significant increase in quality of life. In about 10% of patients the blushing will not be controlled. We do not know why this is the case but it is suspected that in some people the sympathetic nerve fibres controlling blushing of the face bypass the sympathetic chain on their way to the blood vessels in the face. At present there is no known treatment for those unfortunate people in whom the operation does not work. Further research is underway into the reasons why the operation fails and whether it is possible to predict those people on whom it will not work prior to the ETS operation.

Complications.

Complications are unintended outcomes of the operation itself. Examples of complications include BUT ARE NOT LIMITED TO the following:

Death. We are aware of nine deaths worldwide from this procedure. To the best of our knowledge, all of these deaths occurred with surgeons or institutions that were unfamiliar or inexperienced with the ETS procedure.

Bleeding. Bleeding can be a complication of any surgery. The chest cavity, in which the sympathetic chain is situated, contains very large blood vessels which if damaged accidentally during surgery could cause catastrophic bleeding. It is probable that most or all of the deaths reported from ETS surgery were due to bleeding. In experienced hands the risk of major bleeding is extremely low. In our experience of over 500 ETS operations there have been no cases of major bleeding. Results published by the major centres performing ETS surgery confirm that the incidence of severe bleeding is extraordinarily low.

Lung problems. In order to get access to the sympathetic nerves in the chest it is necessary to collapse the lung. At the end of the procedure the lung must be re-expanded and all of the gas removed from the chest cavity. Sometimes the lung does not expand completely and some gas is left inside the chest cavity. This is called a pneumothorax. In most cases this is of no major concern but if the pneumothorax is large a chest tube must be inserted. In even more unusual circumstances there may be a leak of air from the lung requiring the tube to remain in place for one or more days.

A more serious lung problem may be encountered during the surgery if adhesions of the lung to the chest wall are present. Adhesions occur commonly in smokers or in people who have had serious infections of the lung in the past. Often we can deal with these adhesions and still perform the sympathectomy but the risk of bleeding and lung complications after the surgery are increased. These complications can usually be dealt with by a tube into the chest which may need to be present for several days.

On occasions the lung adhesions may be sufficiently severe to cause the operation to be aborted. *The decision to abort the operation is at the surgeon's absolute discretion.* It cannot be predicted preoperatively.

Horner's syndrome. Horner's syndrome is caused by damage to the stellate ganglion. The most obvious sign of Horner's syndrome is a droopy eyelid. Less obvious is constriction of the pupil, a slight sinking of the eyeball and some redness of the white part of the eyes.

The stellate ganglion is part of the sympathetic chain situated above the second rib. With current precise ETS techniques it would be almost impossible to directly damage the stellate ganglion. However any pulling on the nerve or excessive use of electrocautery could damage the stellate ganglion. Damage so caused is extremely uncommon with modern ETS techniques and when it does occur it is usually temporary. Nonetheless, there are recorded instances of Horner's syndrome being permanent. This does not affect eyesight and the droopy eyelid can be fixed by plastic surgery. However, constriction of the pupil, some redness of the whites of the eye and slight recession of the eyeball would be permanent.

Anaesthetic Complications. Modern anaesthesia is extremely safe but in rare instances complications can occur. In some centres it is recommended that only one side of the sympathectomy is done at the one operation, the second being done some weeks later. There have been reports of fluid on the lungs (pulmonary oedema) in association with anaesthesia for operations on both sympathetic chains at the one operation. Our own experience and that of all of the major centres performing sympathectomy is that this complication is so rare that there can be no justification for performing two separate operations to complete the sympathectomy.

Phantom blushing. Following ETS, some patients feel that they are still blushing, but when they look in the mirror there is no blushing. This phenomenon is best explained by the example of patients who have had a limb amputated and still can feel the amputated limb for a while post operatively. The brain has to relearn that there has been a change to the body.

Post-operative Pain. It is common to have some vague pain in the centre of the chest following ETS. A few patients may experience pain along the inner aspect of the arm. This is usually related to trauma to the lower nerve plexus to the arm and in very instances may be permanent.

Nerve Damage. Damage to the nerves between the ribs (intercostal nerves) can occur occasionally. Nerve damage presents as a dull pain or numbness in the inside of the arm or along the route of the rib. In nearly all cases this is temporary and returns to normal within a few weeks or months.

The cause of this is bruising to the nerves during the operation. Very rarely this may be a permanent problem. However it will not affect movement or function of the upper limb.

Other rare complications. There are a myriad of possible complications of any operation about which entire textbooks have been written. It is impossible to detail all possible complications in this document. The prospective patient is again warned that unexpected and unpredictable outcomes can and do occasionally occur after any surgery.

Side-effects Known To Be Associated With Sympathectomy.

Side-effects are direct consequences of the operation other than those intended.

Side-effects can vary from being beneficial to being very serious.

Compensatory or Reflex Sweating. This is the most common and potentially the most serious of the known side-effects of ETS surgery for severe facial blushing. ETS surgery, as well as stopping the blushing, will stop nearly all sweating on the face, scalp and hands. For reasons that we do not fully understand, almost all patients will experience some increased sweating in other areas of the body, principally on the chest, back, groins and thighs. The increased sweating can vary from almost nothing through to an annoyance but in a small percentage of cases, less than 5%, the increased sweating may cause as much or more distress than the original blushing problem.

There is some evidence that compensatory sweating is more likely to be a problem in males, people who are overweight and those who have a tendency to excessive sweating prior to the operation. However there is no sure way to predict who will or who will not have a problem with compensatory sweating.

Compensatory sweating tends to be worse in the first two to three months after the operation and during the first summer after the operation. There is a tendency for it to improve in the 12 to 18 months following surgery but any changes after this are likely to be permanent.

Mild to moderate degrees of compensatory sweating can be improved with the use of anticholinergic drugs such as Ditropan although side-effects such as a dry mouth may limit the effectiveness of these drugs. Botox may be useful if sweating is confined to a particular area that is not too large.

In the unfortunate event of severe compensatory sweating we advocate removal of the clips, preferably in the first 30 to 60 days after the operation is performed. We again emphasise that although there are reports of reversal of compensatory sweating by removal of clips, this has not been confirmed scientifically.

There are also reports of successful reversal of ETS by nerve grafting but likewise there is no scientific proof that nerve grafting can reverse the ETS operation. There are only a very few surgeons in the world who have attempted this procedure and, to our knowledge, none in Australia.

IF YOU PROCEED WITH THE ETS OPERATION FOR SEVERE FACIAL BLUSHING YOU ACCEPT THE POSSIBILITY OF DEVELOPING SEVERE COMPENSATORY SWEATING WHICH MAY NOT BE ABLE TO BE REVERSED.

Gustatory Sweating. Gustatory Sweating is excessive sweating on the temporal region of the face stimulated by the smell or taste of food and drink. This is a very unusual syndrome and the exact cause of this following ETS is not known. It is classed as a side effect of ETS surgery. It occurs in less than 10% of patients undergoing ETS for facial blushing and is usually a minor irritation.

Sometimes an application of Robinul gel to the affected area prior to meal times may be of help. Botox may also help.

If the gustatory sweating is severe then removal of the clips may be necessary. Once again there is no scientific evidence that this will reverse the side effect but there is some anecdotal evidence that it is worth trying in severe cases.

Hands. Your hands will be completely dry after the ETS procedure. This can sometimes cause difficulties with handling paper and may require regular use of moisturising creams or lotions. Initially your hands will be very warm but this effect usually wears off after about six months and some people paradoxically develop cold hands at a later date.

Changes to the Heart and Circulation. Sympathectomy causes changes to the heart and circulation which are well documented. The heart rate is slowed but the heart compensates by pumping slightly more blood with each heart beat. The maximum heart rate in exercise is decreased but for most people there is no change in physical capacity. There is anecdotal evidence that very high performing athletes may suffer a small decrease in maximum physical capacity and if you are an elite athlete you should discuss this very carefully with your surgeon.

Minor changes to blood pressure can occur but these tend, if of any significance at all, to be of more benefit than harm.

Very rare reports have been made of recurrent episodes of fainting after ETS. Whilst probably related to the sympathectomy, the exact cause is not known.

Sweating of the Feet. We know of no reason why sympathectomy at the T2 level should affect foot sweating. Nonetheless, some people report increased sweating, some report decreased sweating and others see no change in sweating of the feet.

Prickling Sensations on Scalp. Some patients notice an unpleasant prickling sensation on the scalp after eating, particularly spicy meals.

Side-effects Reported but Not Known to Be Related to Sympathectomy.

A huge variety of symptoms have been ascribed to the sympathectomy by patients, particularly those who have not had a good result or who have other side-effects of the procedure.

These include symptoms such as loss of libido, tiredness, hair loss, funny feeling in the chest, pains in various parts of the body and many other unexpected symptoms.

Whilst we cannot entirely exclude that these symptoms may be due to the sympathectomy, we believe that most of these symptoms are incidental or psychological as there is no known physiological relationship between the sympathectomy and the symptoms reported.

Summary.

ETS surgery for facial blushing must always be considered a last resort and only undertaken when the blushing is causing severely effecting quality of life. For the majority of patients who undertake the operation there will be a significant improvement in quality of life and for man patients a completely new life and existence. A few people will unfortunately not get any benefit from the operation. In undergoing the operation you accept a small risk that you will end up with side-effects which may be as severe as or even more severe than the original blushing problem.

Complications, however uncommon, may also occur and these are an inherent risk of surgery.

Chris Hensman, Ray McHenry & Richard Gilhome.

May 2008